
When analyzing glycans, Fluorescent derivatization method is often used before measurement by HPLC. Although large amounts of unreacted derivatizing reagents remain in the solution immediately after the derivatization reaction, only unreacted reagents can be removed by performing solid-phase extractions prior to injecting them into the HPLC.


Using MonoSpin, a spin-type solid-state column, separate sugar chains derivatized with 2aminobenzamide (2-Aminobenzamide,2-AB) from unreacted reagents. As a result of evaluating the two types of MonoSpin Amide and MonoSpin NH2, both of them recover sugar chains of more than trisaccharide.

(Y. Yui, C. Aoyama, S. Ota)



# MonoSpin

The MonoSpin series is a spin-column using silica monoliths with uniform continuum pores. Silica monoliths with high porosity are used as carriers, so they can be passed through by centrifugal manipulation alone. Therefore, it is possible to purify and concentrate samples by a simple operation in a short time. It is also suitable when the sample volume is small because the bed volume is small and the liquid can be cut off.









Enlarged view of

The procedure is centrifugation.

Silica monolith disk


# Purification using MonoSpin





### If MpnoSpin NH2 is Used Instead of MonoSpin Amide

MonoSpin NH2 works not only in the HILIC mode but also in the negative ion-exchange mode, but the effects of this study were almost equivalent to those of MonoSpin Amide.



GL Sciences disclaims any and all responsibility for any injury or damage which may be caused by this data directly or indirectly. We reserve the right to amend this information or data at any time and without any prior announcement.

GL Sciences, Inc. Japan 22-1 Nishishinjuku 6-Chome Shinjuku-ku, Tokyo, 163-1130, Japan Phone: +81-3-5323-6620 Fax: +81-3-5323-6621 Email: world@gls.co.jp Web: www.glsciences.com

#### ku, Tokyo, 5652 AS Eindhoven , Japan The Netherlands

The Netherlands Phone: +31 (0)40 254 95 31 Email: <u>info@glsciences.eu</u> Web: www.glsciences.eu

**GL Sciences B.V.** 

**De Sleutel 9** 

### **GL Sciences (ShangHai) Ltd.**

Tower B, Room 2003, Far East International Plaza, NO,317 Xianxia Road, Changning District. Shanghai, China P.C. 200032 Phone: +86 (0)21-6278-2272 Email: <u>contact@glsciences.com.cn</u> Web: www.glsciences.com.cn

### **GL Sciences, Inc. USA**

4733 Torrance Blvd. Suite 255 Torrance, CA 90503 Phone: 310-265-4424 Fax: 310-265-4425 Email: info@glsciencesinc.com Web: www.glsciencesinc.com



International Distributors Visit our Website at:

https://www.glsciences.com/company/distributor.html